

You may also experience some dizziness, blurred vision or short-term memory loss, but these will pass quickly. If you experience any worrying after effects, you should contact your anaesthetist.

● GOING HOME

Make sure there is someone to accompany you home. For the next 24 hours do not:

- Drive a car/motorcycle
- Make important decisions
- Use any dangerous equipment or tools
- Sign any legal documents
- Drink alcohol or take sedatives

● RISKS FROM GENERAL ANAESTHESIA

Modern anaesthesia is very safe. There are however risks and side effects. While most are minor and temporary, some can be serious and long term. Certain conditions increase the risks/complications such as heart and kidney disease, diabetes, high blood pressure, being overweight, colds/flu, and smoking. Complications are also more likely if you are having a major or an emergency surgery.

Common side effects: Feeling sick and vomiting, headache, dizziness, shivering, sore/dry throat, pain and bruising at injection sites.

Less common side effects: Muscle aches, weakness, itching or rash.

Uncommon side effects: Awareness under general anaesthesia, damage to teeth/dental prosthesis, temporary damage to the voice, allergic reactions, damage to nerves.

Very uncommon side effects: Severe allergy causing heart failure, very high temperature, heart attack or stroke, paralysis, vomit in the lungs (causing pneumonia), blood clots in the lungs, brain injury or

● RISKS OF REGIONAL/LOCAL ANAESTHESIA

Reactions or allergies to local anaesthetics are extremely rare. Regional anaesthesia very rarely causes nerve damage due to infection, bleeding and inflammation. This can cause weakness and/or numbness that is usually temporary.

There are specific risks with epidural and spinal anaesthesia. These are:


Headache: The risk is about 1 in 100; the headache is severe, although temporary and can be treated if it occurs.

Backache: Usually temporary and due to bruising from the injection.

Nerve damage: This is extremely rare, but can range from mild, temporary numbness to severe and permanent paralysis.

We encourage you to ask questions. Your anaesthetist will be happy to answer them and discuss the best way to work with you for the best possible outcome.

ANAESTHESIA DURING YOUR SURGERY



**ALPHA INTERNATIONAL WOMEN'S SPECIALISTS
(SINGAPORE) PTE LTD**

(UEN: 201837990C)

101 Irrawaddy Road #12-12

Royal Square At Novena, Singapore 329565

Tel: +65 6860 8888 Fax: +65 6860 8889

© Copyright 2019. All Rights Reserved. Version November 2020.



**ALPHA IVF CENTRE
ALPHA WOMEN'S SPECIALISTS**

Anaesthesia plays a major role to ensure your comfort and safety during your procedure. The following information will help prepare you to receive anaesthetic and know what to expect when you wake up after surgery.

● YOUR ANAESTHETIST

Anaesthetists are doctors with specialised training who:

- Discuss types of anaesthesia and risks with you
- Plan your anaesthetic and pain control
- Are responsible for administering your anaesthesia, monitor your well-being and safety throughout the procedure.
- Make your experience as pleasant and pain free as possible

● TYPES OF ANAESTHESIA

- **General Anaesthesia**
- **Regional Anaesthesia**
- **Local Anaesthesia**
- **Sedation**

The type of anaesthesia you will receive depends on the type of surgery, medical conditions you may have and the length of the operation. Your anaesthetist will discuss the options with you.

Sometimes your anaesthetist will use a combination of anaesthetics. It depends on what is best for you.

● ON THE DAY OF YOUR OPERATION

Nil by mouth = Nothing to eat or drink

The doctors or nurses will advise you how long to fast. It is important to follow these instructions. Food or fluid in the stomach may enter your lungs while you are unconscious. If you don't follow this rule of fasting, the operation may be postponed in the interests of your safety.

If you are taking medications, you should continue to take them, e.g. high blood pressure or asthmatic medications, unless your anaesthetist or surgeon tells you not to.

If you feel unwell on the day that you are coming into Alpha IVF Centre for an elective operation, please call and inform your doctor.

● BEFORE YOUR OPERATION

Your anaesthetist will talk to you before your operation to check on your health, perform relevant examinations and discuss your anaesthetic in detail. They may also send you for blood or other tests. The anaesthetist would need to know:

- 1 If you have a history of high blood pressure, heart problems, thyroid problems, diabetes, asthma, history of reflux or heartburn, or any other medical conditions.
- 2 The past operations you have had and if there were any problems with anaesthesia.
- 3 If you had any abnormal reactions to any drugs or if you have any allergies.
- 4 If you are currently taking any medications. If you are taking aspirin, non-steroidal anti-inflammatory agents or other blood thinning drugs, consult your surgeon or anaesthetist on whether you should stop taking them before your surgery. Please bring with you all your current medications in their original packaging.
- 5 If you have any loose or capped teeth, crowns, or wear dentures.
- 6 If you smoke. Giving up smoking for several weeks before the operation reduces the risks of breathing problems. If you cannot stop smoking, cutting down on smoking will help.

● THE ANAESTHETIC PROCEDURE

General anaesthesia

You get an injection or inhale a gas that makes you unconscious. A tube may be placed in your windpipe to help you breathe. This is essential for some operations. Your anaesthetist stays with you throughout the operation to ensure that you are receiving the correct amount of anaesthetic. There are 2 ways of starting a general anaesthesia:

- Anaesthesia drugs are injected into your body through a cannula placed in a vein
- You will be asked to breathe in oxygen or anaesthesia gases through a mask

Once you are unconscious, anaesthesia can be maintained with intravenous drugs or a mixture of gases, which you will breathe. While you remain unaware of what is happening around you, the anaesthetist monitors your condition closely and constantly adjusts the level of anaesthesia.

As soon as the operation is completed, the drugs will be stopped or reversed so that you regain consciousness.

Regional anaesthesia

This involves injections to numb the nerves around the area of the body where the surgeon needs to operate. This may make the area numb for several hours. You can remain awake, or your anaesthetist may give you an injection to make you relaxed and drowsy. Examples of regional anaesthetics include epidurals for labour and spinal anaesthesia for caesarean section or vaginal hysterectomy.

Local anaesthesia

This involves injections that numb only the site of operation. You will be awake but feel no pain.

Sedation

The anaesthetist administers drugs to make you relaxed and drowsy. This is often used for gastroscopy, colonoscopy, some plastic surgery and IVF procedures.

● AFTER THE OPERATION

You will be observed after surgery to ensure your recovery is as smooth and trouble-free as possible. Your anaesthetist, assisted by the recovery room staff will continue to monitor your condition carefully.

You will feel drowsy for a little while after you wake up from the anaesthetic. You may have a sore or dry throat, feel sick or have a headache. These are temporary and you will recover from them soon.

To help the recovery process, you may be given oxygen to breathe, usually by a clear plastic face mask, and encouraged to take deep breaths and to cough.

Only when you're fully awake and comfortable will you be transferred back to your room or waiting area before returning home.